సేశమవారం 30 అక్టోబర్ 2023

ఇండియన్ నేషనల్ డిఫెన్స్ యూనివర్సిటీ ఎక్కడ ఉంది?

ఇండియన్ మిలిటరీ అకాడమీ ఎక్కడ ఉంది?

డెహాడూన్

కాలేజ్ ఆఫ్ డిఫెన్స్ మేనేజ్మెంట్ ఎక్కడ ఉంది? మిలిటరీ కాలేజ్ ఆఫ్ ఎల్మక్టానిక్స్ అండ్ మెకానికల్ ఇంజినీరింగ్ ఎక్కడ ఉంది?

సికిందాబాద్ సికింబాబాద్ (తిరుమలగిల)

The temporary hardness of water is due to the ..

23 అక్టోబర్ తరువాయి

- Metallic or interstitial hydrides: The transition elements and rare earth metals combine with hydro gen to give interstitial hyd rides. They exhibit metallic properties and are powerful reducing agents. They are non stoichiometric com pounds and their composition varies with temperature and pressure.
- eg. $LaH_{2.76}$, $TiH_{1.73}$ 4. Polymeric hydrides: These are solids containing mole cu les, linked together in two or three dimensions by hydro gen bridge bonds. e.g.: (Be H_2 _n, (MgH_2) _n and (AlH3)_n

HYDROGEN PEROXIDE

Discovery: French chemist The nard 1818.

Occurrence: Traces in air, rain and plants.

PREPARATION OF HYDROGEN PEROXIDE

1. Lab method: From true peroxide by the action of ice cold dil. H2SO4

 $Na_2O_2 + H_2SO_4 \rightarrow Na_2SO_4 + H_2O_2$ $BaO_2 + H_2SO_4 \rightarrow BaSO_4 + H_2O_2$

(HNO₃ is not used since it will oxidise H_2O_2).

2. Merck process: By passing CO2 through a suspension of $Ba\bar{O}_2$ in ice cold water.

 $BaO_2 + H_2O + CO_2 \rightarrow BaCO_3 + H_2O_2$

MANUFACTURE OF HYDROGEN PEROXIDE

1. By electrolysis of 50% ice cold H₂SO₄

2 H 2 S O 4 = H 2 + H 2 S 2 O 8

 $H_2S_2O_8+H_2O\rightarrow H_2SO_4+H_2SO_5$


 $H_2SO_5 + H_2O \rightarrow H_2SO_4 + H_2O_2$

Mechanism:

 $H_2SO_4 \rightleftharpoons H^+ + HSO_4^-$ At anode : ${}^2HSO_4^- \rightarrow H_2S_2O_8 + 2e^-$ At cathode : $^2H^+ + 2e^- \rightarrow H_2$

2. Auto oxidation: Most recent method used in America. In this method the anthraqui none is reduced to anthraqu -inol by dissolving in an organic solvent and passing hyd rogen in presence of Pd. On frothing, the anthraquinol derivative with air, 20% solution of H₂O₂ is obtained and anthraquinone is regenerated.





3. By electrolysis of ammonium sulphate solution and sulp huric acid. When aqueous solution of ammonium sulp hate and sulphuric acid in equimolar proportion is elect -rolysed at low temperature ammonium persulphate is formed. The latter on distillation with sulphuric acid gives 30% solution of hydrogen peroxide.

 $(NH_4)_2SO_4 + H_2SO_4 \rightarrow 2NH_4.HSO_4$ N H 4 H S O 4 ← N H 4 S O 4 + H +

anode $2 NH_4SO_4^- \rightarrow (NH_4)_2 S_2O_8 + 2e^-$ At cathode : ${}^2H^+ + 2e^- \rightarrow H_2(g)$ $(NH_4)_2 S_2 O_8 + 2H_2 O \rightarrow 2 NH_4 H S O_4 + H_2 O_2$

CONCENTRATION OF HYDROGEN PEROXIDE

• It is very carefully concent rated since it decomposes on heating or on standing. $2H_2O_2 \rightarrow 2H_2O + O_2$ (auto-oxidation)

Decomposition is catalysed by Cu, Ag, Pt, Co, Fe, MnO₂ etc. The methods employed for concentration are

- 1. Evaporation: By careful evaporation of solution on a water bath (50% H_2O_2 is obtained).
- 2. Dehydration in vacuum **desicator**: The $50\% H_2O_2$ is dehydrated in a vacuum desi -cator in presence of conc. H_2SO_4 when 90% H_2O_2 is obtained.
- 3. Vacuum distillation: The 90% H₂O₂ obtained in step (ii) is distilled under reduced pressure to get 100% H₂O₂
- **4.** Cooling: The traces of water left are removed by freezing in a freezing mixture when

crystals of hydrogen peroxide separate out.

STRENGTH OF HYDROGEN PEROXIDE

The strength of hydrogen pero xide is indicated in terms of the volume of oxygen at NTP that 1 volume of H₂O₂ gives on heating. For example "20 volume H₂O₂ means 1 volume of H_2O_2 at NTP will give 20 volume of oxygen. The normality and percentage strength of H₂O₂ can be calculated as follows

 $2H_2O_2 \rightarrow 2H_2O + O_2$

 $2 \times 34 = 68 \text{ g}$ 1 mole $O_2 = 22.4$ litres of O_2 at NTP

22.4 litres of O2 at NTP are evolved from 68g of H₂O₂

x litres of O2 at NTP would be evolved

from $\frac{68}{22.4}$ × x g of H₂O₂

where x is volume strength of H₂O₂ Hence strength of x volume of H2O2 =

where 17 is the equivalent weight of H₂O₂

STORAGE OF HYDROGEN PEROXIDE

• It is stored in presence of traces of alcohol, acetanilide or sodium pyrophosphate which slow down the rate of decomposition of hydrogen peroxide.

CHEMICAL PROPERTIES OF HYDROGEN PEROXIDE

1. Acidic nature: It is weakly acidic in nature and pure hydrogen peroxide turns blue litmus into red. (Ka = $1.57 \times$ 10-12 at 293 K). It ionises in two steps H₂O₂ H++

 $H_2O_2 \stackrel{\rightleftharpoons}{=} H^+ + \frac{HO_2^-}{2}$ $HO_{2}^{-} \rightleftharpoons H^{+} + O_{2}^{2-}$

Hence it forms two series of salts eg. NaHO₂ sodium hydroperoxide and Na₂O₂

IIT/NEET **Foundation CHEMISTRY**

(Sodium peroxide)

Oxidising agent: It is strong oxidising agent in acidic as well as in basic medium.

 $H_2O_2 + 2H^+ + 2e^- \rightarrow 2H_2O$ E° = 1.77 V $H_2O_2 + OH^- + 2e^- \rightarrow 3OH^-$ E° = 0.88 V $H_2O_2 \rightarrow H_2O + O$ $PbS + 4O \rightarrow PbSO_4$

 $2FeSO_4 + H_2SO_4 + O \rightarrow Fe_2(SO_4)_1 + H_2O$

 $[Fe(CN)_6] + H_2O + O \rightarrow 2K_3[Fe(CN)_6] + 2KOH$ $\begin{array}{ccc} H_2Cr_2O_7 + 4O \rightarrow & 2CrO_5 & + H_2O \\ Chromic & Blue peroxide \\ acid & of chromium \end{array}$ $C_6H_6 + O \rightarrow C_6H_5OH$

In basic medium

 $2\mathrm{Cr}^{3+} + 3\mathrm{H}_2\mathrm{O}_2 + 10\mathrm{OH}^- \to 2\mathrm{CrO}_4^{2-} + 8\mathrm{H}_2\mathrm{O}$ $\mathrm{Mn^{2+}} + \mathrm{H_2O_2} + \mathrm{2OH}^- \rightarrow \mathrm{MnO_2} + \mathrm{2H_2O}$

In basic medium

 $2Cr^{3+} + 3H_2O_2 + 10OH^- \rightarrow 2CrO_4^{2-} + 8H_2O$ $\mathrm{Mn^{2+}} + \mathrm{H_2O_2} + \mathrm{2OH}^- \rightarrow \mathrm{MnO_2} + \mathrm{2H_2O}$

3.Reducing agent :

a.In acidic medium

 $2MnO_4^- + 6H^+ + 5H_2O_2 \rightarrow 2Mn^{2+} + 8H_2O + 5O_2$

 $Cr_2O_7^{2-} + 8H^+ + 3H_2O_2 \rightarrow 2Cr^{3+} + 7H_2O + 3O_2$ b.In basic medium

 $2K.[Fe(CN),]+2KOH+H.O. \rightarrow 2K.[Fe(CN),]+2H.O+O.$

4.Bleaching properties: Its bleaching action is due to oxidation reaction. $H,O, \rightarrow H,O+O$

 $dye + O \rightarrow dye$ is oxidised and bleached

STRUCTURE OF PEROXIDE

It is represented as follows





HYDROGEN

TEST OF HYDROGEN PEROXIDE

1.It liberates I₂ from acidified KI $2KI + H_2O_2 \rightarrow 2KOH + I_2$

2.Perchromic acid: $H_2Cr_2O_7 + 4H_2O_2 \rightarrow 2CrO_5 + 5H_2O$ $(acidified\ H_2O_2 + amyl\ alcohol + K_2Cr_2O_2) \rightarrow blue\ colour$

3. With titanic sulphate it gives orange red pertitanic acid

 $Ti(SO_4)_2 + H_2O_2 + 2H_2O \rightarrow 2H_2SO_4 + H_2TiO_4$ 4.Black lead sulphide turned white $PbS + 4H_2O_2 \rightarrow PbSO_4 + 4H_2O$

USES OF HYDROGEN PEROXIDE

It is used as a bleaching agent, disinfectant, source of power (90% H₂O₂ as fuel in submarines, rockets and helicopters), in restoration of old paintings in which lead oxide is used as white paint.

WATER

Water is one of the most abundant substances in nature. The 4/5th of the earth surface is covered with water.

SOURCES OF WATER

The sources of water are

1.Surface water

a. Flowing water - streams and rivers

b. Still water - ponds, lakes and reservoirs

2.Underground water - water from wells

3. Rain water 4.Sea water

TYPES OF IMPURITIES PRESENT IN WATER

Dissolved impurities

Inorganic salts eg. : Ca²⁺, Mg²⁺, Fe²⁺, Al³⁺, Na⁺, K⁺ traces of Zn²⁺ and Cu²⁺ (cations) and Cl-, F- etc. (anions) Gases eg.: CO₂, N₂, O₂, oxides of nitro gen, H₂S etc.Organic salts Suspended impurities:

Inorganic: eg.: sand and clay

Organic: eg.: animal matter, vegetable etc.

Colloidal impurities: Finely divided clay, Al(OH)3, Fe (OH)3 colouring matter etc.

Bacterial impurities: Micro-organisms and bacteria

Effect of impurities: The impu -rities effect the followings

- Colour
- Taste
- Hardness
- Alkalinity
- Turbidity Odour

SOFT AND HARD WATER

The water which produces large amount of lather with soap is known as soft water and which forms a scum with soap is known as hard water.

TYPES OF HARDNESS OF WATER

- Temporary hardness: It is due presence of the bicarbonates of calcium or magnesium or both.
- Permanent hardness: It is due to the presence of chlorides and sulphates of calcium and magnesium.

K. Bharathi Co-founder Ed-tech for IIT/NEET foundation

Ph:8309335876

